

Bladder problems in women: childbirth, menopause and aging

Bladder symptoms in women can change after childbirth, around menopause and with aging, but common does not mean inevitable or untreatable. Describing when leakage occurs and whether you empty well is the first step, and treatment should fit the cause and what you want to improve.



Figure 32.1. The Female Bladder Across the Lifespan. Urinary function changes through interacting life stages rather than one inevitable aging pathway.

Key points

- Female urinary symptoms reflect more than the bladder: tissues, childbirth, pelvic support, hormones, bowels, mobility, nerves and medications all matter.
- Stress leakage, urgency leakage, overflow and difficulty reaching the toilet have different causes, and more than one can occur together.
- Delivery history is context, not destiny.
- In later life, getting to the toilet can matter as much as bladder capacity.
- Treatment follows mechanism, severity, risk and preference, not age or embarrassment.

Get prompt care if you notice

- Blood in the urine
- Recurrent infections
- Pain, or difficulty emptying the bladder
- New neurologic findings

Questions to ask your urologist

- What type of bladder or pelvic-support problem do I have?
- Is my bladder emptying well?
- Which life-stage factors, such as childbirth, menopause, medications or mobility, may be contributing?
- What is the least burdensome treatment that fits the mechanism you've identified?
- Would pelvic-floor therapy help, and should it focus on strengthening or relaxation?
- Which findings would make further testing, such as cystoscopy or urodynamics, worthwhile?

Find a urologist near you: [localurology.com](https://www.localurology.com) · Adapted from *A Urology Consult in a Book* by David Shusterman, MD, board-certified urologist. General information, not medical advice. Talk with your own doctor about your situation.